



1. All children should be fully vaccinated for the nine vaccine preventable diseases before the age of one year.
2. Unvaccinated children have great chance to exposed for disease, disability and even for death
3. Vaccinate your children based on the vaccination appointment and health education given by health professional
4. Let a newborn take Polio and BCG immediately after birth
5. Put immunization card in a safe place.
6. Don't forgot vaccination dates, take the immunization cards with you when you go to vaccination posts.
7. All Community participation is mandatory for the achievement of quality and better vaccination coverage
8. Vaccinate, save life, prevent disability.
9. Get your child vaccinated; and ensure his/her good health and bright future
10. Vaccination is the right of children and must for parents
11. Vaccination services are available at any government health facilities free of charge.
12. Take off your responsibility by contributing for the completion of vaccination for defaulted children
13. All under- five children should get polio vaccination during campaigns despite their immunization status.
14. Children may face temporary fever or pain following to vaccination
15. There is no health problems associated with repeated dose given for a child who has vaccinated in routine immunization program.
16. All children should be vaccinated during the campaign irrespective of prior regular vaccination history

17. Take any one with acute flaccid paralysis of one or both of the extremities to the nearby health facility.
18. Prevent polio by improving personal and environmental hygiene and vaccination
19. Wash your hands with soap or ash after using latrines in order to be prevented from polio and other communicable disease
20. Use family latrines properly and be protected from polio virus and other contagious diseases
21. Children living in refugee camps should be vaccinated for polio either through routine immunization or campaign
22. Children with sign and symptoms of AFP should be immediately taken to the nearest health facilities for better management
23. If you observe a child with fever and rash, immediately report to the nearby health facility to prevent possible measles sickness.
24. For an infant < 28 days of age who is unable to suck should be immediately taken to the nearest health facilities for better management of neonatal tetanus/NNT
25. If an infant born healthy and died within 28 days of age, it should be immediately reported to the nearest health facilities